

A Gauge Mediation Model of Dynamical Supersymmetry Breaking without Color Instability

Yasunori Nomura^{a)}, K. Tobe^{b)}, and T. Yanagida^{a)}

a) *Department of Physics, University of Tokyo,
Bunkyo-ku, Hongou, Tokyo 113, Japan*

b) *Department of Physics, The Ohio State University,
Columbus, Ohio 43210, USA*

Abstract

We construct a gauge mediation model of dynamical supersymmetry breaking (DSB) based on a vector-like gauge theory, in which there is a unique color-preserving true vacuum. The DSB scale $\Lambda/4\pi$ turns out to be as high as $\Lambda/4\pi \simeq 10^{8-9}$ GeV, since the transmission of the DSB effects to the standard model sector is completed through much higher loops. This model is perfectly natural and phenomenologically consistent. We also stress that the dangerous D-term problem for the messenger $U(1)_m$ is automatically solved by a charge conjugation symmetry in the vector-like gauge theory.

In the last couple of years there has been a growing interest in dynamical supersymmetry breaking (DSB) with gauge mediation [1], since it provides natural solutions to various phenomenological problems such as flavor-changing neutral current problem and large CP violation. Although many realistic models for gauge mediation have been constructed [2, 3, 4, 5, 6, 7], no satisfactory and simple model has been found yet.

Among various models proposed so far the original model with messenger quarks and leptons in Ref. [2, 3] seems the most attractive and natural. However, this original model has color-breaking global minima [8] and hence one needs to postulate that our universe sits at a local minimum preserving the color SU(3).

The purpose of this letter is to improve the original model so that the true vacuum indeed preserves the color SU(3). In the present model we keep all attractive features in the original model [2, 3]. The present model is perfectly natural and phenomenologically consistent. We also point out that a serious D-term problem for the messenger gauge interaction is automatically solved by a natural symmetry exists in the present model.

The model is based on DSB in vector-like gauge theories observed by Izawa and one of the authors (T.Y.) [9] and Intriligator and Thomas [10]. We adopt here a supersymmetric (SUSY) SU(2) gauge theory with four doublet chiral superfields Q_i and six singlet ones $Z^{ij} = -Z^{ji}$, where i and j are flavor indices ($i, j = 1, \dots, 4$). This theory has a flavor SU(4)_F symmetry. We introduce the following tree-level superpotential to break explicitly the unwanted global SU(4)_F ;

$$W_{\text{tree}} = \lambda_{ij} Z^{ij} (Q_i Q_j). \quad (1)$$

To make our point clear we impose a global SP(4)_F and then the superpotential is written in a simpler form as

$$W_{\text{tree}} = \lambda Z (QQ) + \lambda_Z Z^a (QQ)_a, \quad (2)$$

where (QQ) and Z are singlets of SP(4)_F and $(QQ)_a$ and Z^a are five-dimensional representations of SP(4)_F.

The low energy effective superpotential may be written [11] in terms of gauge-invariant low-energy degrees of freedom

$$V \sim \frac{4\pi}{\Lambda} (QQ), \quad V_a \sim \frac{4\pi}{\Lambda} (QQ)_a \quad (3)$$

as follows:

$$W_{\text{eff}} = X(V^2 + V_a^2 - \frac{\Lambda^2}{(4\pi)^2}) + \frac{\lambda}{4\pi}\Lambda ZV + \frac{\lambda_Z}{4\pi}\Lambda Z^a V_a, \quad (4)$$

where X is an additional chiral superfield and Λ a dynamical scale of the $\text{SU}(2)$ gauge interaction. Factors of 4π are added by using a naïve dimensional analysis (NDA) [12]. For a relatively large value of the coupling λ_Z , we obtain the $\text{SP}(4)_F$ invariant condensation,

$$\langle V \rangle = \frac{\Lambda}{4\pi}, \quad (5)$$

which implies

$$\langle (QQ) \rangle \equiv \langle \frac{1}{2}(Q_1 Q_2 + Q_3 Q_4) \rangle \sim (\frac{\Lambda}{4\pi})^2. \quad (6)$$

Then, the low-energy effective superpotential is approximately given by

$$W_{\text{eff}} \simeq \frac{\lambda}{(4\pi)^2} \Lambda^2 Z. \quad (7)$$

We have nonvanishing F term, $\langle F_Z \rangle \simeq \lambda \Lambda^2 / (4\pi)^2 \neq 0$, and hence SUSY is broken [9, 10].

We gauge a $\text{U}(1)_m$ subgroup in the $\text{SP}(4)_F$.¹ The charge assignments of chiral superfields under $\text{U}(1)_m$ are assumed to be

$$\begin{aligned} & Q_1(+1), \quad Q_2(-1), \quad Q_3(0), \quad Q_4(0), \\ & Z^{12}(0), \quad Z^{34}(0), \quad Z^{13}(-1), \quad Z^{14}(-1), \quad Z^{23}(+1), \quad Z^{24}(+1), \end{aligned} \quad (8)$$

where the numbers in each parentheses denote the $\text{U}(1)_m$ charges. Note that the above condensation Eq. (5) does not break the $\text{U}(1)_m$ gauge symmetry, since V is neutral under the $\text{U}(1)_m$.²

Here, we comment on the light degrees of freedom in the DSB sector in the present model. The present model has a larger global symmetry $\text{SP}(4)_F \times \text{U}(1)_\chi \times \text{U}(1)_R$ and the $\text{U}(1)_\chi$ is spontaneously broken by the condensation of Eq. (6). However, there appears no massless Nambu-Goldstone boson, since the $\text{U}(1)_\chi$ has $\text{SU}(2)$ gauge anomaly. The

¹ Gauging the $\text{U}(1)_m$ symmetry does not restore SUSY.

² We assume the minimal Kähler potential for the Z field for simplicity. Z may get a non-zero vacuum expectation value if one includes higher-order corrections to the Kähler potential, since the Z direction is flat at the tree level. Even if it is the case, however, the $\text{U}(1)_m$ is not broken since Z has also vanishing charge of the $\text{U}(1)_m$, and our main conclusion is unchanged.

't Hooft's anomaly matching condition for the remaining $\text{SP}(4)_F$ and the nonanomalous $\text{U}(1)_R$ is satisfied by the Z field. Thus, the chiral superfield Z is only the flat direction in the present model. This fact is a crucial feature different from the original model [2, 3] which contains light fields with nonvanishing $\text{U}(1)_m$ charges in the DSB sector.

We now consider the messenger sector. The messenger fields are nothing but those in Ref. [2, 3] : we have three chiral superfields,

$$E(+1), \quad \bar{E}(-1), \quad S(0), \quad (9)$$

and vector-like messenger quark and lepton superfields, $d, \bar{d}, l, \bar{l}$. Here, the messenger quark multiplets $d, \bar{d}$ and lepton multiplets $l, \bar{l}$ are all neutral under $\text{U}(1)_m$. We assume that the d and $\bar{d}$ transform as the right-handed down quark and its antiparticle under the standard model (SM) gauge group, respectively. The l and $\bar{l}$ are assumed to transform as the left-handed lepton doublet and its antiparticle, respectively. The most general superpotential for the messenger sector without any dimensional parameters is

$$W_{\text{mess}} = k_E S E \bar{E} + \frac{f}{3} S^3 + k_d S d \bar{d} + k_l S l \bar{l}. \quad (10)$$

This is the same form as in the original model [2, 3] and natural in the sense that there is a consistent symmetry which forbids other terms in the superpotential.

When we switch on the $\text{U}(1)_m$ gauge interaction, the soft SUSY-breaking masses for E and $\bar{E}$ fields are generated through the SUSY-breaking effects in the DSB sector. The resulting scalar potential is

$$V_{\text{mess}} = \sum_{\eta} \left| \frac{\partial W}{\partial \eta} \right|^2 + m_E^2 |E|^2 + m_{\bar{E}}^2 |\bar{E}|^2, \quad (11)$$

where m_E and $m_{\bar{E}}$ are soft SUSY-breaking masses for E and $\bar{E}$, and η denotes chiral superfields $E, \bar{E}, S, d, \bar{d}, l$, and $\bar{l}$. One of the essential differences between the present model and the original one is that the soft SUSY-breaking mass squared m_E^2 and $m_{\bar{E}}^2$ are positive in the present model while these are negative in the original model. We will discuss this point below.

There are two types of quantum corrections to the soft SUSY-breaking masses for the messenger fields: one is from 'threshold effects' and the other from 'running effects'. In

Ref. [2, 3], the m_E^2 and $m_{\bar{E}}^2$ come essentially from the 'running effects'. The DSB sector contains light superfields $\chi^\pm$ in the original model which have $U(1)_m$ charges. Since both of SUSY-breaking soft masses and the SUSY-invariant mass for the superfields $\chi^\pm$ are very small compared with the DSB scale Λ , the 'running effects' from the SUSY-breaking scale to the χ field mass scale becomes important. According to the two-loop renormalization group equation (RGE), we find that the m_E^2 and $m_{\bar{E}}^2$ receive negative contributions from the χ and gauge multiplet loops, which results in negative m_E^2 and $m_{\bar{E}}^2$. With these negative SUSY-breaking masses the DSB effects are transmitted to the messenger quarks and leptons, which causes, however, instability of the color-preserving vacua [8].

On the other hand, the 'threshold effects' dominate over the 'running effects' in the present model. The reason is that since the DSB sector is a vector-like theory and the flavor $SP(4)_F$ is unbroken, all composite fields have dynamical masses of order Λ and there are no light field besides the Z superfield as discussed above. Therefore, the 'running effects' which would drive the m_E^2 and $m_{\bar{E}}^2$ negative is negligible compared with relatively large 'threshold effects'. As a consequence we have positive SUSY-breaking masses for the E and $\bar{E}$ fields.

Let us evaluate the threshold corrections to the m_E^2 and $m_{\bar{E}}^2$. It may be very difficult to calculate them directly, because of the strong gauge dynamics in the SUSY breaking sector. Therefore, we estimate them by using the effective theory resulting from the strong gauge dynamics. The effective theory contains many composite fields which consist of the elementary Q fields. Here, we assume that the lightest composite fields are a pair of chiral superfields Φ and $\bar{\Phi}$ which have $U(1)_m$ charges ± 1 , respectively.³ We also consider that they give a dominant contribution to the masses m_E^2 and $m_{\bar{E}}^2$.

Following the NDA prescription in Ref. [12], the effective theory concerning Φ , $\bar{\Phi}$, Z fields is described by

$$\mathcal{L} = \frac{\Lambda^4}{g_2^2} \left[\int \frac{d^4\theta}{\Lambda^2} \tilde{K}(\phi, \phi^\dagger, \bar{\phi}, \bar{\phi}^\dagger) + \int \frac{d^2\theta}{\Lambda} \tilde{W}(\phi, \bar{\phi}, z) + \text{h.c.} \right], \quad (12)$$

where g_2 is a strong $SU(2)$ gauge coupling ($g_2 \simeq 4\pi$) and $\tilde{W}$ is a function of dimensionless

³ In fact, it has been shown that the lightest bound state of the SUSY QED consists of a pair of chiral supermultiplets in Ref. [13]. This fact may support our assumption.

superfields $\phi = g_2\Phi/\Lambda$, $\bar{\phi} = g_2\bar{\Phi}/\Lambda$, and $z = \lambda Z/\Lambda$ as

$$\tilde{W} \simeq \phi\bar{\phi} + z\phi\bar{\phi}, \quad (13)$$

where we have assumed that all coupling constants of Eq.(13) are $O(1)$. This gives us the following effective superpotential:⁴

$$W_{\text{comp}} \simeq \Lambda\Phi\bar{\Phi} + \lambda Z\Phi\bar{\Phi}. \quad (14)$$

Then, the m_E^2 and $m_{\bar{E}}^2$ can be evaluated perturbatively as long as $\lambda \lesssim O(1)$. The two-loop diagrams relevant to these masses are shown in Fig. 1. The resulting m_E^2 and $m_{\bar{E}}^2$ are positive and their order of magnitudes are given by

$$m_E = m_{\bar{E}} \sim \frac{\alpha_m}{4\pi} \frac{\lambda F_Z}{\Lambda} \simeq \frac{\alpha_m}{4\pi} \frac{\lambda^2}{16\pi^2} \Lambda, \quad (15)$$

where $\alpha_m = g_m^2/4\pi$ and g_m is the $U(1)_m$ gauge coupling constant. We note that this result is also obtained by applying directly the NDA to Q -integration without the introduction of the composite chiral superfields.

Unlike in the original model [2, 3], the SUSY-breaking effects are not transmitted to the messenger quarks and leptons with these positive soft SUSY-breaking masses. Thus, the squarks, sleptons, and gauginos in the minimal supersymmetric SM (MSSM) sector remain massless at this level.

Now, we discuss transmission of the DSB effects to the MSSM sector. For the positive m_E^2 and $m_{\bar{E}}^2$, the S and F_S do not develop vacuum-expectation values. In the present model, however, the S field gets a negative soft SUSY-breaking mass squared $-m_S^2$ from the 'running effects' through loops of the E and $\bar{E}$ fields with the Yukawa coupling $k_E S E \bar{E}$ in Eq. (10).⁵ The m_S^2 is evaluated as

$$m_S^2 \simeq \frac{4}{(4\pi)^2} k_E^2 m_E^2 \ln \frac{\Lambda}{m_E}. \quad (16)$$

Altogether, the scalar potential for the messenger sector becomes

$$V_{\text{mess}} = |k_E E \bar{E} + f S^2 + k_d d \bar{d} + k_l l \bar{l}|^2$$

⁴ For the Kähler potential, we assume the minimal form $\tilde{K} = \phi^\dagger \phi + \bar{\phi}^\dagger \bar{\phi}$, which leads to $K_{\text{comp}} = \Phi^\dagger \Phi + \bar{\Phi}^\dagger \bar{\Phi}$.

⁵ This possibility has been also noted in the original model [2].




Figure 1: Diagrams contributing to masses of the scalar components of E and $\bar{E}$ chiral supermultiplets. Here, g and $\tilde{g}$ represent $U(1)_m$ gauge boson and gaugino, respectively. Scalar and fermion components of the chiral supermultiplets are denoted by the same symbols of the chiral superfields and those with tilde, respectively.

$$\begin{aligned}
& +|k_E S E|^2 + |k_E S \bar{E}|^2 \\
& +|k_d S d|^2 + |k_d S \bar{d}|^2 \\
& +|k_l S l|^2 + |k_l S \bar{l}|^2 \\
& +m_E^2 |E|^2 + m_E^2 |\bar{E}|^2 - m_S^2 |S|^2.
\end{aligned} \tag{17}$$

This potential has a global minimum at

$$\begin{aligned}
\langle S^* S \rangle &= \frac{m_S^2}{2f^2}, \\
\langle E \rangle &= \langle \bar{E} \rangle = \langle d \rangle = \langle \bar{d} \rangle = \langle l \rangle = \langle \bar{l} \rangle = 0, \\
\langle |F_S| \rangle &= \frac{m_S^2}{2f},
\end{aligned} \tag{18}$$

in a certain parameter region. Thus, all the SM gauge symmetries are preserved and the SUSY-breaking effects are transmitted to the messenger quark and lepton multiplets through $\langle F_S \rangle$.⁶

When we integrate out the messenger quarks and leptons d , $\bar{d}$, l , and $\bar{l}$, the MSSM gauginos acquire masses at the one-loop level as

$$m_{\tilde{g}_i} = c_i \frac{\alpha_i}{4\pi} \Lambda_{\text{mess}}, \tag{19}$$

where $c_1 = 5/3$, $c_2 = c_3 = 1$, and $m_{\tilde{g}_i}$ ($i = 1, \dots, 3$) denote the bino, wino, and gluino masses, respectively. The soft SUSY-breaking masses for squarks, sleptons, and Higgses $\tilde{f}$ in the MSSM sector are generated at the two-loop level as

$$m_{\tilde{f}}^2 = 2\Lambda_{\text{mess}}^2 \left[C_3 \left(\frac{\alpha_3}{4\pi} \right)^2 + C_2 \left(\frac{\alpha_2}{4\pi} \right)^2 + \frac{5}{3} Y^2 \left(\frac{\alpha_1}{4\pi} \right)^2 \right], \tag{20}$$

where $C_3 = 4/3$ for color triplets and zero for singlets, $C_2 = 3/4$ for weak doublets and zero for singlets, and Y is the SM hypercharge ($Y = Q_{\text{em}} - T_3$). Here, Λ_{mess} is an effective messenger scale defined as

$$\Lambda_{\text{mess}} \equiv \frac{\langle |F_S| \rangle}{\langle S \rangle} = \frac{m_S}{\sqrt{2}}, \tag{21}$$

⁶ The messenger gauge group $U(1)_m$ is also unbroken in this vacuum. Although the massless $U(1)_m$ gauge boson couples to the SM quarks and leptons through higher loops, we find that this causes no phenomenological problem, since induced forces are short range and very weak. We should note also that there are stable $U(1)_m$ charged particles in the present model. All such particles, however, have the masses of order $\Lambda \simeq 10^{9-10}$ GeV. Thus, the cosmological density of these particles is expected to be very low since the reheating temperature T_R of the inflation should be $T_R \lesssim 10^6$ GeV [14] to solve the gravitino problem in the present model, and hence they cause no phenomenological problem.

which can be written in terms of the SUSY-breaking scale $\sqrt{F_Z}$ as

$$\begin{aligned}\Lambda_{\text{mess}} &\simeq \frac{\sqrt{2}}{(4\pi)^4} \alpha_m \lambda^2 k_E \sqrt{\ln \frac{(4\pi)^3}{\alpha_m \lambda^2}} \cdot \Lambda \\ &= \frac{\sqrt{2}}{(4\pi)^4} \alpha_m \lambda \sqrt{\lambda} k_E \sqrt{\ln \frac{(4\pi)^3}{\alpha_m \lambda^2}} \cdot \sqrt{F_Z}.\end{aligned}\quad (22)$$

If we intend to make the MSSM gaugino and sfermion masses at the electroweak scale, the effective messenger scale Λ_{mess} must be 10^{4-5} GeV. Then, the SUSY-breaking scale $\sqrt{F_Z}$ becomes 10^{8-9} GeV for $\alpha_m \simeq 10^{-2}$ and $\lambda \sim k_E \sim O(1)$.⁷ This corresponds to the gravitino mass,

$$m_{3/2} \simeq \frac{F_Z}{\sqrt{3} M_G} \simeq 10 \text{ MeV} - 1 \text{ GeV}, \quad (23)$$

where M_G is the gravitational scale $M_G \simeq 2.4 \times 10^{18}$ GeV. Thus, dangerous flavor-changing neutral currents are suppressed well below the present experimental limits [16].

Finally, we should stress that the dangerous D-term problem [2] for the $U(1)_m$ is automatically solved in the present model. That is, there is a charge conjugation symmetry in the DSB sector defined as

$$\begin{aligned}Q_1 &\leftrightarrow Q_2, \quad V \leftrightarrow -V, \\ Z^{12} &\leftrightarrow -Z^{12}, \quad Z^{13} \leftrightarrow Z^{23}, \quad Z^{14} \leftrightarrow Z^{24},\end{aligned}\quad (24)$$

which forbids the Fayet-Iliopoulos D-term of V [3, 17]. Here, V is the $U(1)_m$ gauge superfield. This may be a big merit of the present vector-like gauge theory [9, 10].

In conclusion, we have constructed a gauge mediation model which has a unique color-preserving global minimum in a certain parameter region.⁸ The SUSY-breaking scale turns out to be as high as $\langle \sqrt{F_Z} \rangle \sim 10^{8-9}$ GeV which results in $m_{3/2} \simeq 10 \text{ MeV} - 1 \text{ GeV}$. The present model is perfectly natural and phenomenologically consistent. However, it should be noted that if there exist moduli fields like a dilaton in superstring theories with masses $\sim m_{3/2}$, the present model has a serious cosmological problem, since decays

⁷ In order that the $U(1)_m$ gauge coupling constant should not blow up below the grand unification scale $\simeq 2 \times 10^{16}$ GeV, α_m at the Λ_{mess} must be smaller than $\sim 3 \times 10^{-2}$ [15]. A similar requirement for the Yukawa coupling constants λ and k_E leads to $\lambda \lesssim 2$ and $k_E \lesssim 1$ at the Λ_{mess} .

⁸ It may not be excluded that the original model [2, 3] has also such a desired global minimum in the nonperturbative regime.

of such moduli fields produce too much background γ rays as pointed out in the recent work [18]. However, if the presence of light moduli fields is an inevitable prediction of the superstring theories is not yet clear to us.

One of the authors (K.T.) would like to thank T. Moroi for useful comments. The work of K.T. is supported by DOE contract DOE/ER/01545-726.

References

- [1] M. Dine, W. Fischler, and M. Srednicki, *Nucl. Phys.* **B189** (1981) 575;
S. Dimopoulos and S. Raby, *Nucl. Phys.* **B192** (1981) 353;
L. Alvarez-Gaumé, M. Claudson, and M. Wise, *Nucl. Phys.* **B207** (1982) 96;
C. Nappi and B. Ovrut, *Phys. Lett.* **B113** (1982) 175.
- [2] M. Dine, A.E. Nelson, and Y. Shirman, *Phys. Rev.* **D51** (1995) 1362.
- [3] M. Dine, A.E. Nelson, Y. Nir, and Y. Shirman, *Phys. Rev.* **D53** (1996) 2658.
- [4] T. Hotta, K.-I. Izawa, and T. Yanagida, *Phys. Rev.* **D55** (1997) 415.
- [5] K.-I. Izawa, Y. Nomura, K. Tobe, and T. Yanagida, *Phys. Rev.* **D56** (1997) 2886;
Y. Nomura and K. Tobe, hep-ph/9708377.
- [6] E. Poppitz and S. Trivedi, *Phys. Rev.* **D55** (1997) 5508;
N. Arkani-Hamed, J. March-Russell, and H. Murayama, hep-ph/9701286;
H. Murayama, *Phys. Rev. Lett.* **79** (1997) 18;
S. Dimopoulos, G. Dvali, G. Giudice, and R. Rattazzi, hep-ph/9705307;
M. Luty, hep-ph/9706554;
Y. Shirman, hep-ph/9709383.
- [7] L. Randall, *Nucl. Phys.* **B495** (1997) 37;
N. Haba, N. Maru, and T. Matsuoka, *Nucl. Phys.* **B497** (1997) 31; *Phys. Rev.* **D56** (1997) 4207;
S. Raby, *Phys. Rev.* **D56** (1997) 2852;
Y. Shadmi, *Phys. Lett.* **B405** (1997) 99;
K.-I. Izawa, *Prog. Theor. Phys.* **98** (1997) 443;
C. Csaki, L. Randall, W. Skiba, hep-ph/9707386.
- [8] I. Dasgupta, B.A. Dobrescu, and L. Randall, *Nucl. Phys.* **B483** (1997) 95;
N. Arkani-Hamed, C.D. Carone, L.J. Hall, H. Murayama, *Phys. Rev.* **D54** (1996) 7032.
- [9] K.-I. Izawa and T. Yanagida, *Prog. Theor. Phys.* **95** (1996) 829.
- [10] K. Intriligator and S. Thomas, *Nucl. Phys.* **B473** (1996) 121.

- [11] For a review, K. Intriligator and N. Seiberg, *Nucl. Phys. Proc. Suppl.* **45BC** (1996) 1.
- [12] M.A. Luty, hep-ph/9706235;
A.G. Cohen, D.B. Kaplan, and A.E. Nelson, *Phys. Lett.* **B412** (1997) 301.
- [13] W. Buchmuller, S.T. Love, and R.D. Peccei, *Nucl. Phys.* **B204** (1982) 429.
- [14] T. Moroi, H. Murayama, and M. Yamaguchi, *Phys. Lett.* **B303** (1993) 289.
- [15] A. Gouvêa, T. Moroi, and H. Murayama, *Phys. Rev.* **D56** (1997) 1281.
- [16] F. Gabbiani, E. Gabrielli, A. Masiero, and L. Silvestrini, *Nucl. Phys.* **B477** (1996) 321.
- [17] W. Fishler, H.P. Nilles, J. Polchinski, S. Raby, and L. Susskind, *Phys. Rev. Lett.* **47** (1981) 757.
- [18] M. Kawasaki and T. Yanagida, *Phys. Lett.* **B399** (1997) 45;
J. Hashiba, M. Kawasaki, and T. Yanagida, hep-ph/9708226 to appear in *Phys. Rev. Lett.*